Mondstuk van die Suid-Afrikaanse aartappelbedryf • Mouthpiece of the South African potato industry

CHIPS

VOL 35 NO 03 • MAY / JUNE 2021

MARKMONITOR: DIE EERSTE DRIE MAANDE VAN 2021 OP VARSPRODUKTEMARKTE

POTATO TUBER MOTH INSECTICIDES, MODES OF ACTION, AND SENSIBLE USE

Mechanisation training for Limpopo producers

PSA's roll-out of its mass media campaign

Limpopo-kultivarproef onder besproeiing: Dendron

Potato tuber moth insecticides, their modes of action, and sensible use to prevent or delay resistance development

Principal author: Desireé van Heerden • Contributors: Dr Fienie Niederwiesser, Roleen la Grange and Dr Gerhard Verdoorn

Insecticides are grouped according to an international system developed by the Insecticide Resistance Action Committee (IRAC). The IRAC Mode of Action (MoA) classification groups each active ingredient according to a specific chemical class and resistance group, based on the MoA of the active ingredient.

Insecticides and their IRAC grouping

The IRAC classification guides the use of different modes of action successively to prevent sensitivity shift, and to prevent or delay the development of insecticide resistance (visit *www.irac-online.org* or download the IRAC app for smart devices).

During the past few seasons, higher than normal potato tuber moth (PTM) populations were reported and the dry and hot seasons most likely contributed to these numbers. Warm winters with high temperatures during the day and little to no frost allow earlier than normal occurrence of PTM. Seasons with high moth 'explosions' are not an unknown phenomenon and occur every few years, especially during droughts and warm periods.

Insect life cycles are temperature dependent and are normally shortened by increased temperatures, resulting in increased numbers of PTM generations in one season. A higher percentage of unmarketable potato tubers (>30%) was harvested during the past three years, compared with the usual expected 5 to 7%.

Insecticides are classified according to five different categories based on physiological functions that are affected by insects:

- *Growth:* Insect development is mainly controlled by juvenile hormones, by directly perturbing cuticle formation/deposition or lipid biosynthesis. Insect growth regulators are generally slow- to moderately slow-acting.
- *Midgut: Lepidopteran*-specific microbes or their derived microbial toxins that are sprayed or expressed in transgenic crop varieties (not applicable to potatoes at this stage). Moderately acting.
- *Respiration:* Several insecticides interfere with mitochondrial respiration by the inhibition of electron transport and/or oxidative phosphorylation. Generally fast- to moderately fast-acting.
- *Nerve and muscle function:* Most current insecticides act on nerve and muscle targets. Insecticides that act on these targets are generally fast-acting.
- *Unclassified functions:* Several insecticides are known to affect less well-described target sites or functions, or to act non-specifically on multiple targets.

By far the largest class of insecticides affects the nerve and muscle system of insects. This collective class consists of pyrethroids, carbamates, organophosphates, avermectins, and spinosyns, among others, each with a different MoA and specific target site.

Figure 1: The different target sites for controlling PTM.



Table 1: Active ingredients that are registered for the control of PTM in potatoes.

IRAC group	Chemical class	Active ingredient	MoAs/physiological functions affected
1A	Carbamate	Methomyl	Acetylcholinesterase (AChE) inhibitors Nerve action
1B	Organophosphates	Acephate, azinphos-methyl, methamidophos, methidathion and profenofos	AChE inhibitors Nerve action
3A	Pyrethroids	Alpha-cypermethrin, beta-cyfluthrin, bifenthrin, deltamethrin, esfenvalerate, gamma-cyhalothrin and lambda-cyhalothrin	Sodium channel modulators Nerve action
4A and 3A	Neonicotinoids and pyrethroids	Acetamiprid and bifenthrin	Nicotinic acetylcholine receptor (nAChR) competitive modulators Nerve action
4A and 15	Neonicotinoids and benzoylureas	Acetamiprid and novaluron	nAChR competitive modulators, and chitin synthesis inhibitors Nerve action and growth regulation
5	Spinosyns	Spinetoram and spinosad	nAChR allosteric modulators – Site 1 Nerve action
13	Pyrroles	Chlorfenapyr	Uncouplers of oxidative phosphorylation via disruption of the proton gradient Energy metabolism
14	Nereistoxin analogue	Cartap hydrochloride	nAChR channel blockers Nerve action
15	Benzoylureas	Diflubenzuron, lufenuron and novaluron	Inhibitors of chitin biosynthesis affecting CHS1 Growth regulation
22A	Oxadiazines	Indoxacarb	Voltage-dependent sodium channel blockers Nerve action
28	Diamides	Chlorantraniliprole and cyantraniliprole	Ryanodine receptor modulators Nerve and muscle action
28 and 3A	Diamides and 3A	Chlorantraniliprole and lambda-cyhalothrin	Ryanodine receptor and sodium channel modulators Nerve and muscle action
UN	Pyridalyl	Pyridalyl dichloropropene derivative	Compounds of unknown or uncertain MoAs
UNF	Fungus	Beauveria bassiana	Compounds of unknown or uncertain MoAs
Pheromone and 3A	Pheromone and pyrethroids	(E,Z)-4,7 & (E,Z,Z)-4,7,10-tridecatrienyl acetate and permethrin	Lure and sodium channel modulators Lure and nerve action

Carbamates and organophosphates are both Group 1 but are categorised into Sub-group 1A and Sub-group 1B, respectively. To prevent sensitivity shift and the development of insecticide resistance, chemical classes of different MoAs (resistance groups) should be rotated, and consecutive insect generations should not continuously be exposed to the same MoAs (*Figure 1*).

Twelve different IRAC insecticide classes corresponding to their equivalent resistance groups are registered in South Africa for the control of the PTM, *Phthorimaea operculella* (*Table 1*). Study the table and plan accordingly to develop a spray programme that will prevent sensitivity shifts and resistance development.

Performance of active ingredients

Not all the active ingredients in a specific chemical class share the same characteristics. Plant uptake can be by contact, translaminar, or systemic action, while insect uptake can be through stomach or contact action.

The formulation type and quality, such as EC, SC, WP, rain fastness, stability in sunlight, mixability and compatibility with other products, application type and method, all play a crucial role in the efficacy of an active ingredient. Choosing an active ingredient requires taking all these characteristics into account along with the plant size, level of pest

infestation and the pest complex present, to select the best option.

Take note again that active ingredients within the same IRAC group share the same MoAs. Carbamates and organophosphates, for example, are both in Group 1 and share the same MoAs. A few examples of active ingredients registered against PTM are mentioned below. Always read the entire label to ensure that the products are applied correctly.

Carbamates (Group 1A)

- *Methomyl:* Contact and stomach action. Efficacy greatly reduced after spray residues have dried. Rapidly degrades in soil with a short half-life (DT_{50} 4 to 8 days) at 20°C. Soil moisture and pH are important in the breakdown of methomyl, which is not compatible with alkaline products. Relatively stable in sunlight.

Organophosphates (Group 1B)

- *Acephate:* Contact and stomach action. Moderate persistence. Residual activity of 10 to 21 days. Non-phytotoxic. Readily biodegraded and non-persistent in soil with a short half-life (DT_{50} 2 days).
- *Azinphos-methyl:* Non-systemic. Contact and stomach action. Rapidly hydrolysed in alkaline and acidic media. Half-life of several weeks under normal conditions (DT_{50} 87 days at pH 4 and 50 days at pH 7; only 4 days at pH 9). Low soil mobility. Photodegradation on soil surface.
- *Profenofos:* Non-systemic. Contact and stomach action. Exhibits translaminar effect and ovicidal properties. Not compatible with sulphur, alkaline products or captan. Do not mix with metal-containing compounds or apply with nitrogen-containing foliar feeds. Short half-life in soil (DT_{50} 1 week).
- *Methamidophos:* Systemic. Contact and stomach action. Rapidly degraded in soil. Short half-life in soil (DT_{50} <2 days). Photolysis contributes to rapid degradation.

Pyrethroids (Group 3A)

- *Bifenthrin:* Non-systemic. Contact and stomach action. Not compatible with alkaline products. Long half-life in soil (DT_{50} 65 days).
- *Lambda-cyhalothrin:* Non-systemic. Contact and stomach action, with repellent properties and rapid knockdown effect. Stable to light, stable in storage (>6 months), but rapidly degraded in soil, especially under dry conditions (DT_{50} 4 weeks). Strongly adsorbed to soil. Does not leach.

Neonicotinoids (Group 4A)

- *Acetamiprid:* Systemic, with translaminar movement in the plant. Stomach and contact activity. High potential for bioaccumulation. Mobile in soil, but degrades rapidly. Low potential for leaching into groundwater. Half-life between <1 and 8.2 days.

Spinosyns (Group 5)

- *Spinosad:* Non-systemic. Contact and stomach action. Short soil half-life (DT_{50} 9 to 17 days). Low pH value (< 6) of the spray mixture will decrease the residual performance. Performs best at pH 6 to 9.
- *Spinetoram:* Non-systemic. Contact and stomach action. Rapidly degrades in soil (DT_{50} 3 to 5 days).

Pyrroles (Group 13)

- *Chlorfenapyr:* Limited systemic activity. Mainly stomach with some contact action. Pro-insecticide (metabolised into an active insecticide after entering the host). Persistent in soil but binds very strongly to soil particles and does not leach.

Nereistoxin analogues (Group 14)

- *Cartap hydrochloride:* Systemic, with stomach and contact action. Stable in acidic conditions but hydrolyses in neutral or alkaline solution. Short soil half-life (DT_{50} 3 days).

Figure 2: Tuber infested with PTM due to being exposed under leaves and not being completely covered.

Benzoylureas (Group 15)

- *Lufenuron:* Insect growth regulator. Non-systemic. Translaminar effect, with strong stomach and moderate contact activity. Larvicidal, ovicidal and transovarial action. Reduces egg fecundity. Stable at pH 5 to 7 and a very long half-life (DT_{50} 512 days). Strong adsorption onto soil particles. Not compatible with carbamates or alkaline products.

Oxadiazines: (Group 22A)

- *Indoxacarb:* Contact and stomach action. Rapidly terminates insect feeding on crop. Moderately persistent but immobile in soil (DT_{50} 3 to 23 days).

Figure 4: Sealed ridges.

Diamides (Group 28)

- *Cyantraniliprole*: Systemic effect by soil uptake with some translaminar movement. Active through ingestion and contact. Ovicidal, ovi-larvicidal and adulticide activity. Low soil mobility due to high soil adsorption, with moderate photodegradation and a short half-life (DT_{50} 4 to 25 days). Degrades rapidly. Resistance risks seem higher than other MoAs.
- *Chlorantraniliprole*: Stomach and contact action. Weak translaminar activity. Slow soil degradation (DT_{50} 270 days). Low soil mobility due to high soil adsorption, low water solubility and slow photodegradation. Resistance risks seems higher than other MoAs.

Unclassified (UN)

- *Pyridalyl*: Non-systemic. Contact and stomach action. Moderately slow-acting. Highly immobile in soil with a long half-life (DT_{50} 93 to 182 days).

Effective application and practice

Virtually no insecticide can be drenched into the soil to control PTM after senescence (maturation) and ridging. The best practice is to control it while green plant material is still available for chemical uptake and to minimise exposed tubers (*Figure 2*), soil cracks or soil cavities next to stems (*Figure 3*), especially in shallow bearing cultivars.

Moths can lay eggs close to cracks and young hatching larvae can then move down the cracks to infest tubers. Make sure cracks are sealed by irrigation after ridging (*Figure 4*).

Potato tubers that are slightly exposed are very easy targets for larvae; field observations showed that the tubers that were infested with PTM larvae were the ones very close to the soil surface with some level of exposure. It is therefore strongly advised to minimise exposed tubers to the absolute minimum.

Figure 3: Soil cracks and cavities.

This article is Monograph 1 of the Potato Production Stewardship Programme, a collaborative initiative of Potatoes South Africa, CropLife South Africa and the IRAC. For more information, contact Dr Gerhard H Verdoorn at Gerhard@CropLife.co.za.